arXiv:1406.6269v1 [math.AT] 24 Jun 2014

RATIONAL HOMOTOPY THEORY OF FUNCTION SPACES AND HOCHSCHILD COHOMOLOGY

ILIAS AMRANI

Abstract. Given a map $f : X \to Y$ of simply connected spaces of finite type such. The space of based loops at f of the space of maps between X and Y is denoted by $\Omega_f \mathrm{Map}(X, Y)$. For $n > 0$, we give a model categorical interpretation of the existence (in functorial way) of an injective map of $\mathbb{Q}$-vector spaces $\pi_n \Omega_f \mathrm{Map}(X, Y_{\mathbb{Q}}) \to \mathrm{HH}_k^{-n}(C^*(Y), C^*(X)_f)$, where HH_k^* is the (negative) Hochschild cohomology and $C^*(X)_f$ is the rational cochain complex associated to X equipped with a structure of $C^*(Y)$-differential graded bimodule via the induced map of differential graded algebras $f^* : C^*(Y) \to C^*(X)$. Moreover, we identifiy the image in presice way by using the Hodge filtration on Hochschild cohomology. In particular, when $X = Y$, we describe the fundamental group of the identity component of the monoid of self equivalence of a (rationalization of) space X i.e., $\pi_1 Aut(X_{\mathbb{Q}})_{id}$.

Introduction

Our main goal in this article is the study of the function space $\mathrm{Map}(X, Y)$ between two **rational** topological spaces from non-commutative point of view. More precisely, for a fixed map $f : X \to Y$ we study the homotopy groups of the path connected component $\mathrm{Map}(X, Y_{\mathbb{Q}})_f$. It is well known [3] that **rationally** (under some finiteness conditions) the homotopy groups $(\pi_n,\ n > 1)$ of $\mathrm{Map}(X, Y_{\mathbb{Q}})_f$ are given by the André-Quillen cohomology $\mathrm{AQ}^{-n}(C^*(Y), C^*(X))$, where $C^*(X)$ is seen as a module over $C^*(Y)$ via the induced map of E_∞-differential graded algebras $f^* : C^*(Y) \to C^*(X)$. The point is that the André-Quillen cohomology is quite complicated to compute. We should notice that we are using the fact that any rational E_∞-differential graded algebra is equivalent to a rational commutative differential graded algebra. Let k be any commutative ring, and denote the model category of E_∞-differential graded k-algebras by $\mathsf{E}_\infty - \mathsf{dgAlg}_k$ and the model category of associative differential graded k-algebras by dgAlg_k. The (derived) forgetful functor $U : \mathsf{E}_\infty - \mathsf{dgAlg}_k \to \mathsf{dgAlg}_k$ induces a map of simplicial sets

$$\alpha : \mathrm{Map}_{\mathsf{E}_\infty - \mathsf{dgAlg}_k}(R, S) \to \mathrm{Map}_{\mathsf{dgAlg}_k}(R, S) := \mathrm{Map}_{\mathsf{dgAlg}_k}(UR, US).$$

In all what follows, we will consider only the positively graded algebras with increasing differentials by degree one. A perfect example is the cochain complex associated to a topological space. The interpretation of the higher homotopy groups is quite

2000 *Mathematics Subject Classification.* Primary 55, Secondary 14 , 16, 18.

Key words and phrases. DGA, CDGA, Mapping Space, Rational Homotopy Theory, Hochschild Cohomology, André-Quillen Cohomology, Harrison Cohomology, Hodge Filtration.

Partially supported by the project CZ.1.07/2.3.00/20.0003 of the Operational Programme Education for Competitiveness of the Ministry of Education, Youth and Sports of the Czech Republic.

simple, in fact in [2], we have shown that for a given map $f : R \to S$ of differential graded k-algebras we have

$$\pi_n \mathrm{Map}_{\mathsf{dgAlg}_k}(R, S)_f \cong \mathrm{HH}_k^{-n+1}(R, S_f) \ \forall \ n > 1, \tag{0.1}$$

where HH_k^* is the Hochschild cohomology and S is seen as R-bimodule via f.

Rational homotopy theory. When $k = \mathbb{Q}$, Sullivan has proven that there is an ∞-equivalence between the category of simply connected rational spaces (finite type) and a subcategory of simply connected commutative differential graded k-algebras (of finite type) [12]. The ∞-equivalence is given by the cochain functor $C^*(-, k)$ after strictification. For any map $f : X \to Y$ of rational simply connected spaces, the forgetful functor U induces the following map of k-vector spaces

$$\pi_{n+1}\alpha : \ \pi_{n+1}\mathrm{Map}(X, Y_{\mathbb{Q}})_f \to \mathrm{HH}_k^{-n}(C^*(Y), C^*(X)_f).$$

In [3, Theorem 3.8], Block and Lazarev give an explicit formula when f is **homotopy equivalent to a constant map**. They (re)proved that (under the convention that the cohomology is negatively graded)

$$\pi_n \mathrm{Map}(X, Y_{\mathbb{Q}})_f \cong \prod_{i=1}^{\infty} \pi_i(Y) \otimes \mathrm{H}^{i-n}(X, \mathbb{Q}).$$

p-Adic homotopy theory. When p is a prime number and $k = \overline{\mathbb{F}}_p$ the algebraic closure of the field with p-elements. Mandell's fundamental theorem [11] says that the ∞-category of p-complete spaces (with some finiteness conditions) is ∞-equivalent to a full ∞-subcategory of E_∞-differential graded k-algebras via the cochain functor $C^*(-, \overline{\mathbb{F}}_p)$. Suppose that $f : X \to Y$ is a map of simply connected spaces (with some finiteness conditions), then the forgetful functor U induces the following map of abelian groups ($n > 0$)

$$\pi_{n+1}\alpha : \ \pi_{n+1}\mathrm{Map}(X, Y_p^\wedge)_f \to \mathrm{HH}_k^{-n}(C^*(Y), C^*(X)_f),$$

where $Y \to Y_p^\wedge$ is a p-completion functor.

Theorem 0.1 (2.2)**.** *Suppose that $k = \mathbb{Q}$. Let $f : X \to Y$ be a map of simply connected spaces of (finite type), then the forgetful functor*

$$U : \mathrm{Map}_{\mathsf{E}_\infty - \mathsf{dgAlg}_k}(C^*(Y), C^*(X)) \to \mathrm{Map}_{\mathsf{dgAlg}_k}(C^*(Y), C^*(X))$$

induces a map of k-vector spaces such that:

1. $[X, Y_{\mathbb{Q}}] = \pi_0 \mathrm{Map}_{\mathsf{E}_\infty - \mathsf{dgAlg}_k}(C^*(Y), C^*(X)) \to \pi_0 \mathrm{Map}_{\mathsf{dgAlg}_k}(C^*(Y), C^*(X))$ *is injective.*
2. $\pi_1 \mathrm{Map}(X, Y_{\mathbb{Q}})_f = \pi_1 \mathrm{Map}_{\mathsf{E}_\infty - \mathsf{dgAlg}_k}(C^*(Y), C^*(X))_f \to \pi_1 \mathrm{Map}_{\mathsf{dgAlg}_k}(C^*(Y), C^*(X))_f$ *is injective map of groups.*
3. $\forall \ n > 0$, *the induced map*

$$\pi_{n+1}\mathrm{Map}(X, Y_{\mathbb{Q}})_f = \pi_{n+1}\mathrm{Map}_{\mathsf{E}_\infty - \mathsf{dgAlg}_k}(C^*(Y), C^*(X))_f \to \mathrm{HH}_k^{-n}(C^*(Y), C^*(X)_f),$$

   is injective map of $\mathbb{Q}$-vector spaces.
4. *If $X = Y$ and $f = id$, then $\pi_1 Aut(X_{\mathbb{Q}})_{id} \to \mathrm{HH}_k^{0,\times}(C^*(X), C^*(X))$ is an injective map of abelian groups.*

The space $X_{\mathbb{Q}}$ is the rationalization of X, $Aut(X)$ is the monoid of self equivalences, and $\mathrm{HH}_k^{0,\times}(C^*(X), C^*(X))$ is the group of invertible elements of the k-algebra $\mathrm{HH}_k^0(C^*(X), C^*(X))$.

Warning 0.2. When $k = \overline{\mathbb{F}}_p$, the induced maps $\pi_n\alpha$ are far to be injective in general.

Theorem 0.3 (Hodge filtration 2.5)**.** *With the same assemption as in precedent Theorem, we have the following isomorphism*

$$\pi_{n+1}\mathrm{Map}(X, Y_{\mathbb{Q}})_f \cong \mathrm{HH}^{-n}_{(1)}(C^*(Y), C^*(X)_f),\ \forall\ n > 0, \forall\ f.$$

1. General framework

For what follows we fix $k = \mathbb{Q}$. Notice that $\mathsf{E}_\infty - \mathsf{dgAlg}_k \simeq \mathsf{dgCAlg}_k$. In the abstract we described only the applications. In order to prove them we pass by the model category of differential graded algebras (commutative and non-commutative). We denote the pointed model category of augmented (resp. commutative and E_∞) differential graded k-algebras by dgAlg^*_k (resp. dgCAlg^*_k and $\mathsf{E}_\infty - \mathsf{dgAlg}^*_k$). Notice that the model structure in the commutative case make sense when k is of characteristic 0. For some technical reasons, we define the functor of cochain complexes $C^*(-, k) = C^*(-) : \mathsf{sSet}^{op} \to \mathsf{E}_\infty - \mathsf{dgAlg}_k$. In this section, a space means a simplicial set.

Notation 1.1. *All differential graded algebras are non-negatively graded and the differentials increase the degree by +1. Consider the map of operads (in the differential graded context)* $\mathsf{Ass} \to \mathsf{Com}$*, since the category of operad is a model category we have the factorization* $\mathsf{Ass} \to \mathsf{E}_\infty \to \mathsf{Com}$*, where the first map is a cofibration and the second map is a trivial fibration. We have shown in* [1, Lemma 1.1]*, that* E_∞ *is admissible and the forgetful functor* $U : \mathsf{E}_\infty - \mathsf{dgAlg}^*_k \to \mathsf{dgAlg}^*_k$ *preserves cofibrant objects and cofibration between cofibrant objects. That is the reason why we work with* E_∞*-operad instead of the operad* Com*.*

Recall that we have a following diagram of (Quillen) adjunctions:

$$\begin{array}{ccc}
\mathsf{dgAlg}_k & \underset{U}{\overset{F}{\rightleftarrows}} & \mathsf{E}_\infty - \mathsf{dgAlg}_k \\
{\scriptstyle -\oplus k}\uparrow \downarrow{\scriptstyle \underline{U}} & & {\scriptstyle -\oplus k}\uparrow \downarrow{\scriptstyle \underline{U}} \\
\mathsf{dgAlg}^*_k & \underset{U}{\overset{F}{\rightleftarrows}} & \mathsf{E}_\infty - \mathsf{dgAlg}^*_k
\end{array}$$

where, F and $\underline{U}$ are left adjoints and U, $\oplus k$ are right adjoints.

Warning 1.2. In what follows, we took the liberty to not denote the forgetful functors i.e., when R is an (augmented) E_∞-differential graded algebra we consider it also as an (augmented) associative differential graded algebra without mentioning the forgetful functor.

Theorem 1.3. [1, Theorem 3.1] *Let $k = \mathbb{Q}$, for any R and S augmented commutative differential graded k-algebras, the forgetful functor $U : \mathsf{E}_\infty - \mathsf{dgAlg}^*_k \to \mathsf{dgAlg}^*_k$ induces a map $\alpha :\ \mathrm{Map}_{\mathsf{E}_\infty - \mathsf{dgAlg}^*_k}(R, S) \to \mathrm{Map}_{\mathsf{dgAlg}^*_k}(R, S)$ such that*

$$\pi_n\mathrm{Map}_{\mathsf{E}_\infty - \mathsf{dgAlg}^*_k}(R, S) \to \pi_n\mathrm{Map}_{\mathsf{dgAlg}^*_k}(R, S)$$

is injective map of groups for $n > 0$. Moreover, the map $\Omega\mathrm{Map}_{\mathsf{E}_\infty - \mathsf{dgAlg}^*_k}(R, S) \to \Omega\mathrm{Map}_{\mathsf{dgAlg}^*_k}(R, S)$ has a functorial retract with respect to the target argument S .

For more details, we refere to 2.2.

Lemma 1.4. *Let k be any field. The (derived) functor $C^*(-) : \mathsf{sSet}^{op} \longrightarrow \mathsf{E}_\infty\mathsf{dgAlg}_k$ commutes with homotopy limits.*

Proof. The functor C^* has a left adjoint (cf [11, Proposition 4.2]), they form a Quillen pair. the homotopy limits in sSet^{op} are the homotopy colimits in sSet, it follows that for any diagram $J \to \mathsf{sSet}$ we have an isomorphism $C^*(\mathrm{hocolim}_{j\in J}X_j) \cong \mathrm{holim}_{j\in J}C^*(X_j)$ in the homotopy category $\mathrm{Ho}(\mathsf{sSet})$. □

Notation 1.5. *We denote the simplicial sphere of dimension n by S^n.*

Definition 1.6. Let R be an augmented E_∞-differential graded k-algebra, we say that R connected if $\pi_0\mathrm{Map}_{\mathsf{dgAlg}_k^*}(R, k \oplus k) = \pi_0\mathrm{Map}_{\mathsf{E}_\infty-\mathsf{dgAlg}_k^*}(R, k \oplus k) = *$.

Lemma 1.7. *Let X be a pointed connected simplicial set, and let $R \in \mathsf{E}_\infty - \mathsf{dgAlg}_k^*$ be connected (cofibrant). Then the induced map by the forgetful functor*

$$\mathrm{Map}_{\mathsf{E}_\infty-\mathsf{dgAlg}_k^*}(R, C^*(X)) \to \mathrm{Map}_{\mathsf{dgAlg}_k^*}(R, C^*(X))$$

has a functorial (depending on X) retract in $\mathrm{Ho}(\mathsf{sSet}_*)$.

Proof. We define two functors $\Psi, \Phi : \mathsf{sSet}_*^{op} \to \mathsf{sSet}_*$ as follows

- $\Psi(X) = \mathrm{Map}_{\mathsf{E}_\infty-\mathsf{dgAlg}_k^*}(R, C^*(X))$ and
- $\Phi(X) = \mathrm{Map}_{\mathsf{dgAlg}_k^*}(R, C^*(X))$.

These functors verify the following properties

(1) They send a weak equivalence $X \to Y$ to a weak equivalence since the functor $C^*(-)$ preserves weak between cofibrant objects and $\mathrm{Map}_{\mathsf{E}_\infty-\mathsf{dgAlg}_k^*}(R, -)$, $\mathrm{Map}_{\mathsf{E}_\infty-\mathsf{dgAlg}_k^*}(R, -)$ preserves weak equivalence between fibrant objects since R is cofibrant as E_∞-algebra and as associative algebra cf 1.1.
(2) The functors Ψ and Φ take homotopy limits to homotopy colimits, it follows that the mapping spaces of a model category commutes with homotopy limits in the second argument and the fact that $C^*(-)$ takes homotopy colimits to homotopy limits 1.4. Moreover the forgetful functor $U : \mathsf{E}_\infty - \mathsf{dgAlg}_k^* \to \mathsf{dgAlg}_k^*$ commutes with homotopy limites.
(3) $\Psi(*)$ and $\Phi(*)$ are contractible since k is a terminal object in $\mathsf{E}_\infty - \mathsf{dgAlg}_k^*$ and dgAlg_k^*.

It follows form [10, Theorem 16], that $\Psi(-)$ and $\Phi(-)$ are representable i.e., there exists two simplicial sets C and A such that $\Psi(-) \simeq \mathrm{Map}_{\mathsf{sSet}_*}(-, C)$ and $\Phi(-) \simeq \mathrm{Map}_{\mathsf{sSet}_*}(-, A)$ in $\mathrm{Ho}(\mathsf{sSet}_*)$ the natural transformation $\Psi(-) \to \Phi(-)$ is represented by a map $C \to A$. By theorem 1.3, we know that the map $\Omega\Psi(-) \to \Omega\Phi(-)$ has a functorial retract (in $\mathrm{Ho}(\mathsf{sSet}_*)$), it follows that the map $\Omega\mathrm{Map}_{\mathsf{sSet}_*}(-, C) \to \Omega\mathrm{Map}_{\mathsf{sSet}_*}(-, A)$ has a functorial retract, it implies that $\Omega C \to \Omega A$ has a retract. On another hand R is connected, it follows that A and C are connected. hence, the induced map $A \to C$ has a retract in $\mathrm{Ho}(\mathsf{sSet}_*)$. We conclude that $\mathrm{Map}_{\mathsf{E}_\infty-\mathsf{dgAlg}_k^*}(R, C^*(X)) \to \mathrm{Map}_{\mathsf{dgAlg}_k^*}(R, C^*(X))$ has a functorial retract in $\mathrm{Ho}(\mathsf{sSet}_*)$ for any simplicial set X. □

Corollary 1.8. *For any connected augmented E_∞-differential graded algebra, and any pointed simplicial set X, the natural map*

$$\mathrm{Map}_{\mathsf{E}_\infty-\mathsf{dgAlg}_k^*}(R, C^*(X)) \to \mathrm{Map}_{\mathsf{dgAlg}_k^*}(R, C^*(X))$$

induces an injective map on homotopy groups.

Theorem 1.9. *Let R be a connected augmented E_∞-differential graded algebra, with augmentation $\nu : R \to k$. Let X be any pointed simplicial set, let $f : R \to C^*(X)$ be any map of augmented E_∞-differential graded algebras. Then the induced map by the forgetful functor*

$$\alpha : \ \mathrm{Map}_{\mathsf{E}_\infty - \mathsf{dgAlg}_k}(R, C^*(X)) \to \mathrm{Map}_{\mathsf{dgAlg}_k}(R, C^*(X))$$

has a functorial retract (on the variable X), in particular $\forall\ f, \forall\ n > 0$:

- $\pi_0\alpha : \pi_0\mathrm{Map}_{\mathsf{E}_\infty - \mathsf{dgAlg}_k}(R, C^*(X)) \to \pi_0\mathrm{Map}_{\mathsf{dgAlg}_k}(R, C^*(X))$ *and*
- $\pi_n\alpha : \pi_n\mathrm{Map}_{\mathsf{E}_\infty - \mathsf{dgAlg}_k}(R, C^*(X))_f \to \pi_n\mathrm{Map}_{\mathsf{dgAlg}_k}(R, C^*(X))_f$

are injective maps

Proof. First of all, notice that we have an obvious cofiber sequence of pointed simplicial sets

$$S^0 \xrightarrow{\ i\ } X_+ \xrightarrow{\ p\ } X$$

where X_+ is the pointed simplicial set $X \coprod *$. It is enough to notice that

$$\mathrm{Map}_{\mathsf{E}_\infty - \mathsf{dgAlg}_k^*}(R, C^*(X_+)) \simeq \mathrm{Map}_{\mathsf{E}_\infty - \mathsf{dgAlg}_k}(R, C^*(X))$$

and

$$\mathrm{Map}_{\mathsf{dgAlg}_k^*}(R, C^*(X_+)) \simeq \mathrm{Map}_{\mathsf{dgAlg}_k}(R, C^*(X)),$$

then the result follows from 1.7.

□

2. Main theorems and applications

Proposition 2.1. Suppose that $k = \mathbb{Q}$. Let R be an augmented E_∞-differential graded k-algebra of finite type (i.e $\dim_k \mathrm{H}^i(R) < \infty\ \forall\ i$) such that $\mathrm{H}^0(R) = k$ and $\mathrm{H}^1(R) = 0$ then R is connected in the sense of 1.6.

Proof. First of all, by adjunction $\mathrm{Map}_{\mathsf{E}_\infty - \mathsf{dgAlg}_k^*}(R, k \oplus k) \simeq \mathrm{Map}_{\mathsf{E}_\infty - \mathsf{dgAlg}_k}(R, k)$. Without loosing generality we can suppose that R is cofibrant as $\mathsf{E}_\infty - \mathsf{dgAlg}_k$, hence R is cofibrant as dgAlg_k (by construction of the operad E_∞ cf 1.1). By Sullivan Theorem $\pi_0\mathrm{Map}_{\mathsf{E}_\infty - \mathsf{dgAlg}_k}(R, k) = *$. It follows that for any maps $\nu : R \to k$ and $\mu : R \to k$ are homotopic in $\mathsf{E}_\infty - \mathsf{dgAlg}_k$. According to [8], we have a commutative diagram in $\mathsf{E}_\infty - \mathsf{dgAlg}_k$



where $P(R)$ is a path object associated to R. Notice that the path object is the same for graded differential associative algebras if we consider $R \in \mathsf{dgAlg}_k$. Since $\mathrm{H}^0(R) = k$ any map $R \to k$ in dgAlg_k is actually a map in $\mathsf{E}_\infty - \mathsf{dgAlg}_k$. We conclude that $\pi_0\mathrm{Map}_{\mathsf{dgAlg}_k}(R, k) = * = \pi_0\mathrm{Map}_{\mathsf{dgAlg}_k^*}(R, k \oplus k)$.

□

Theorem 2.2 (Main Theorem)**.** *Suppose that $k = \mathbb{Q}$. Let $f : X \to Y$ be a map of simply connected spaces of (finite type), then the forgetful functor*

$$U : \mathrm{Map}_{\mathsf{E}_\infty - \mathsf{dgAlg}_k}(C^*(Y), C^*(X)) \to \mathrm{Map}_{\mathsf{dgAlg}_k}(C^*(Y), C^*(X))$$

induces a map of k-vector spaces such that:

1. $[X, Y_{\mathbb{Q}}] = \pi_0 \mathrm{Map}_{\mathsf{E}_\infty - \mathsf{dgAlg}_k}(C^*(Y), C^*(X)) \to \pi_0 \mathrm{Map}_{\mathsf{dgAlg}_k}(C^*(Y), C^*(X))$ *is injective.*
2. $\pi_1 \mathrm{Map}(X, Y_{\mathbb{Q}})_f = \pi_1 \mathrm{Map}_{\mathsf{E}_\infty - \mathsf{dgAlg}_k}(C^*(Y), C^*(X))_f \to \pi_1 \mathrm{Map}_{\mathsf{dgAlg}_k}(C^*(Y), C^*(X))_f$ *is injective.*
3. $\forall\ n > 0$,

$$\pi_{n+1} \mathrm{Map}(X, Y_{\mathbb{Q}})_f = \pi_{n+1} \mathrm{Map}_{\mathsf{E}_\infty - \mathsf{dgAlg}_k}(C^*(Y), C^*(X))_f = \mathrm{AQ}^{-n-1}(C^*(Y), C^*(X)_f)$$

and the induced map

$$\pi_{n+1} \mathrm{Map}(X, Y_{\mathbb{Q}})_f \to \pi_{n+1} \mathrm{Map}_{\mathsf{E}_\infty - \mathsf{dgAlg}_k}(C^*(Y), C^*(X))_f = \mathrm{HH}_k^{-n}(C^*(Y), C^*(X)_f)$$

is an injective map of $\mathbb{Q}$-vector spaces.

4. *If $X = Y$ and $f = id$, then $\pi_1 Aut(X_{\mathbb{Q}})_{id} \to \mathrm{HH}_k^{0,\times}(C^*(X), C^*(X))$ is an injective map of abelian groups.*

Proof. By hypothesis X and Y are of finite type, we deduce by [12] that

$$\mathrm{Map}_{\mathsf{E}_\infty - \mathsf{dgAlg}_k}(C^*(Y), C^*(X))$$

is equivalent to $\mathrm{Map}(X, Y_{\mathbb{Q}})$, on the other hand by Theorem 1.9, the forgetful functor $U : \mathsf{E}_\infty - \mathsf{dgAlg}_k \to \mathsf{dgAlg}_k$ induces an injective map

$$\alpha : \pi_i \mathrm{Map}_{\mathsf{E}_\infty - \mathsf{dgAlg}_k}(C^*(Y), C^*(X))_f \to \pi_i \mathrm{Map}_{\mathsf{dgAlg}_k}(C^*(Y), C^*(X))_f$$

for all $i \geq 0$. Moreover if $i > 1$, Block-Lazarev theorem gives us the isomorphism

$$\pi_i \mathrm{Map}_{\mathsf{E}_\infty - \mathsf{dgAlg}_k}(C^*(Y), C^*(X))_f \cong \mathrm{AQ}^{-i}(C^*(Y), C^*(X)_f),$$

and by [2],

$$\pi_i \mathrm{Map}_{\mathsf{dgAlg}_k}(C^*(Y), C^*(X))_f \cong \mathrm{HH}_k^{-i+1}(C^*(Y), C^*(X)_f)$$

Hence, the induced map α is exactly $\mathrm{AQ}^{-i}(C^*(Y), C^*(X)_f) \to \mathrm{HH}_k^{-i+1}(C^*(Y), C^*(X)_f)$, which is injective map of $\mathbb{Q}$-vector spaces. Applying Sullivan theorem we deduce that $\pi_i \mathrm{Map}(X, Y_{\mathbb{Q}})_f \cong \mathrm{AQ}^{-i}(C^*(Y), C^*(X)_f)$ for $i > 1$. In particular, when $X = Y$ and $f = id$, $\mathrm{Map}(X, X)_{id} = Aut(X)_{id}$ and

$$\mathrm{Map}(X_{\mathbb{Q}}, X_{\mathbb{Q}})_{id} = Aut(X_{\mathbb{Q}})_{id}.$$

Therefore, $\pi_1 Aut(X_{\mathbb{Q}})_{id} \cong \pi_1 \mathrm{Map}_{\mathsf{E}_\infty - \mathsf{dgAlg}_k}(C^*(X), C^*(X))_{id}$. In [2, Corollary 3.6], we have shown that $\pi_1 \mathrm{Map}_{\mathsf{dgAlg}_k}(C^*(X), C^*(X))_{id}$ is isomorphic to the kernel of the natural map of (abelian) groups $\mathrm{HH}_k^{0,\times}(C^*(X), C^*(X)) \to \mathrm{H}^{0,\times}(C^*(X)) = \mathbb{Q}^\times$. The result follows for Theorem 1.9. □

Corollary 2.3. *Let M be a simply connected orientable closed manifold of dimension d, for all $i > 0$, we have an injective map of $\mathbb{Q}$ vector spaces*

$$\pi_i \Omega_{id} Aut(M) \otimes \mathbb{Q} \to \mathrm{H}_{i+d}(\mathcal{L}M, \mathbb{Q}),$$

where $\mathcal{L}M$ is the space of free loops on M, i.e., $\mathrm{Map}(S^1, M)$.

Proof. Since M is a finite CW-complex, it is a direct consequence of Theorem 2.2, and the fact that
$\mathrm{HH}_k^*(C^*(M), C^*(M)) \cong \mathrm{H}_{*+d}(\mathcal{L}M, \mathbb{Q})$ [4]. □

Remark 2.4. Corollary 2.3 was also proven in [5, Theorem 2 (1)] using a different method.

2.1. **Hodge filtration on Hochschild cohomology over a field of characteristic zero.** In our main Theorem 2.2, we have identified the higher homotopy groups of $\mathrm{Map}(X, Y_{\mathbb{Q}})_f$ based at some continuous map $f : X \to Y$ as a sub $\mathbb{Q}$-vector space of the (negative) Hochschild cohomology. According to [6, Theorem 3.1], there exists a Hodge decomposition on the Hochschild cohomology $\mathrm{HH}^*_k(R, S)$ for any differential graded $\mathbb{Q}$-algebra R and any differential graded R-bimodule S. More precisely Ginot has proved in [6], the following formula in the rational case:

$$\mathrm{HH}^*(R, S) \cong \prod_{n\geq 0} \mathrm{HH}^*_{(n)}(R, S),$$

where the $\mathbb{Q}$-vector spaces $\mathrm{HH}^*_{(n)}(R, S)$ are eigenspaces for an iterated power of some operator.

Theorem 2.5. *With the same assemption as in Theorem 2.2, we have the following isomorphism*

$$\pi_{n+1}\mathrm{Map}(X, Y_{\mathbb{Q}})_f \cong \mathrm{HH}^{-n}_{(1)}(C^*(Y), C^*(X)_f),\ \forall\ n > 0, \forall\ f.$$

Proof. First of all, we notice that $\pi_n\mathrm{Map}(X, Y_{\mathbb{Q}})_f \cong \mathrm{AQ}^{-n}(C^*(Y), C^*(X)_f)$ for all $n > 1$ (cf [3]), where AQ^* is the André-Quillen cohomology. On another hand $\mathrm{HH}^{-n}_{(1)}(C^*(Y), C^*(X)_f) = \mathrm{Harr}^{-n}(C^*(Y), C^*(X)_f)$, where Harr^* is the Harrison cohomology, cf [6, Theorem 3.1]. Since we work in characteristic zero, Harrison cohomology and André-Quillen cohomology agree up to a shift, more precisely $\mathrm{AQ}^{n-1} = \mathrm{Harr}^n$. It follows that

$$\begin{aligned}
\pi_{n+1}\mathrm{Map}(X, Y_{\mathbb{Q}})_f &\cong \mathrm{AQ}^{-n-1}(C^*(Y), C^*(X)_f) \\
&\cong \mathrm{Harr}^{-n}(C^*(Y), C^*(X)_f) \\
&\cong \mathrm{HH}^{-n}_{(1)}(C^*(Y), C^*(X)_f),\ \forall\ n > 0, \forall\ f.
\end{aligned}$$

□

Remark 2.6. Theorem 2.5 is a generalization of [5, Theorem 2 (2)].

Appendix

There is a class of model categories called simplicial model categories [7], roughly speaking a simplicial model category is tensored, cotensored and enriched over the model category of simplicial sets in a compatible way (adjunction compatibility, and model structure compatibility). In general a model category C do not need to be simplicial model category. Moreover, a Quillen adjunction between simplicial model categories

$$\mathsf{C} \underset{U}{\overset{F}{\rightleftarrows}} \mathsf{D},$$

is not a simplicial adjunction in general. In [9, Chapter 5, 6], Hovey introduced a notion of module category. We will need a more richer structure and we will call it **enriched module structure**. In the classical context any ordinary category with product and coproduct is an enriched Set-module. More precisely, suppose that D is an ordinary category with products and coproducts, we can define the following functors:

(1) $- \otimes - :\ \mathsf{Set} \times \mathsf{D} \to \mathsf{D}$ such that for any set X and any object $D \in \mathsf{D}$ we have $X \otimes D = \coprod_{i \in X} D$.
(2) $A(-,-) :\ \mathsf{Set}^{op} \times \mathsf{D} \to \mathsf{D}$ such that for any set X and for any $D \in \mathsf{D}$ we define $A(X, D) = \prod_{i \in X} D$.

Definition 2.7. An enriched Set-module D is a category with all products and coproducts such that we have natural isomorphism for any $X, Y \in \mathsf{Set}$ and any $C, D \in \mathsf{D}$

- $(X \times Y) \otimes D \cong X \otimes (Y \otimes D)$.
- $\hom_{\mathsf{D}}(C, A(X, D)) \cong \hom_{\mathsf{D}}(X \otimes C, D)$.
- $\hom_{\mathsf{Set}}(X, \hom_{\mathsf{D}}(C, D)) \cong \hom_{\mathsf{D}}(X \otimes C, D) \cong \hom_{\mathsf{D}}(C, A(X, D))$.
- $* \otimes D \cong D$.

A simplicial category D in the sense of [7] is an enriched sSet-module in the sense of 2.7, where we replace $\hom_{\mathsf{D}}$ by the natural enrichment of D denoted by $\mathrm{Map}_{\mathsf{D}}$ (simplicial set).

Theorem 2.8. *Let* D *be any (pointed) model category, then the homotopy category* $\mathrm{Ho}(\mathsf{D})$ *is an enriched* $\mathrm{Ho}(\mathsf{sSet})$*-module (enriched* $\mathrm{Ho}(\mathsf{sSet}_*)$*-module).*

Proposition 2.9. Given any Quillen adjunction between model categories

$$\mathsf{C} \underset{U}{\overset{F}{\rightleftarrows}} \mathsf{D},$$

it induces the following isomorphisms :

- $\mathrm{Map}_{\mathsf{C}}(X, U(Y)) \cong \mathrm{Map}_{\mathsf{D}}(F(X), Y)$ in $\mathrm{Ho}(\mathsf{sSet})$
- $F(X \otimes C) \cong X \otimes F(C)$ in $\mathrm{Ho}(\mathsf{D})$ for any $X \in \mathsf{sSet}$ and any $C \in \mathsf{C}$.
- $U(A(X, D)) \cong A(X, UD)$ in $\mathrm{Ho}(\mathsf{C})$ for all $X \in \mathsf{sSet}$ and any $D \in \mathsf{D}$

The proof of the precedent theorem and proposition can be deduced from [9]. The involved mapping spaces tensors and cotensors are defined in the **derived** sense, we took the liberty to not specify the derived symbols (e.g. $\mathbf{R}$ and $\mathbf{L}$).

Notation 2.10. *If* D *is a pointed model category, we denote by* ΩD *the object* $A(S^1, D)$ *and* ΣD *the object* $S^1 \otimes D$.

2.2. **Complement to Theorem 1.3.** *We explain, the cited Theorem using the previous language. Let* R *be cofibrant an augmented* E_∞*-differential graded* $\mathbb{Q}$*-algebras. Considering the adjunction*

$$\mathsf{dgAlg}_k^* \underset{U}{\overset{F}{\rightleftarrows}} \mathsf{E}_\infty - \mathsf{dgAlg}_k^*,$$

our theorem says that we have a natural map $S^1 \otimes R \to F(S^1 \otimes UR) \simeq S^1 \otimes FUR$ *which has a retract in* $\mathrm{Ho}(\mathsf{E}_\infty - \mathsf{dgAlg}_k^*)$*. In other words, suppose that* $S \in \mathsf{E}_\infty - \mathsf{dgAlg}_k^*$*, we have a retract in* $\mathrm{Ho}(\mathsf{sSet}_*)$ *of the map*

$$\mathrm{Map}_{\mathsf{E}_\infty - \mathsf{dgAlg}_k^*}(S^1 \otimes R, S) \to \mathrm{Map}_{\mathsf{E}_\infty - \mathsf{dgAlg}_k^*}(S^1 \otimes FUR, S)$$

which can be rewritten by using adjunctions as:

$$h : \Omega\mathrm{Map}_{\mathsf{E}_\infty - \mathsf{dgAlg}_k^*}(R, S) \to \Omega\mathrm{Map}_{\mathsf{dgAlg}_k^*}(UR, US),$$

such that, there is an induced left inverse map r*, i.e.,* $r \circ h = id$ *and it is functorial with respect to* S.

References

[1] Ilias Amrani. Comparing commutative and associative unbounded differential graded algebras over $\mathbb{Q}$ from homotopical point of view. *arXiv preprint math/1401.7285*, 2013.

[2] Ilias Amrani. The mapping space of unbounded differential graded algebras. *arXiv preprint arXiv:1303.6895*, 2013.

[3] Jonathan Block and Andrej Lazarev. André–Quillen cohomology and rational homotopy of function spaces. *Advances in Mathematics*, 193(1):18–39, 2005.

[4] Ralph L Cohen and John DS Jones. A homotopy theoretic realization of string topology. *Mathematische Annalen*, 324(4):773–798, 2002.

[5] Yves Félix and Jean-Claude Thomas. Monoid of self-equivalences and free loop spaces. *Proceedings of the American Mathematical Society*, 132(1):305–312, 2004.

[6] Grégory Ginot. On the Hochschild and Harrison (co)homology of C_∞-algebras and applications to string topology. In *Deformation Spaces*, pages 1–51. Springer, 2010.

[7] P.G. Goerss and JF Jardine. *Simplicial homotopy theory*. Birkhauser, 1999.

[8] Vladimir Hinich. Homological algebra of homotopy algebras. *Communications in algebra*, 25(10):3291–3323, 1997.

[9] M. Hovey. Model categories. *Mathematical Surveys and Monographs*, 63, 1999.

[10] JF Jardine. Representability theorems for simplicial presheaves. *Preprint*, 2009.

[11] Michael A Mandell. E_∞-algebras and p-adic homotopy theory. *Topology*, 40(1):43–94, 2001.

[12] Dennis Sullivan. Infinitesimal computations in topology. *Publications Mathématiques de l'IHÉS*, 47(1):269–331, 1977.

Department of Mathematics, Masaryk University, Kotlarska 2, Brno, Czech Republic.

E-mail address: ilias.amranifedotov@gmail.com

E-mail address: amrani@math.muni.cz